



30+ YEARS OF  
PATIENT  
SUPPORT\*

**RealSupport**<sup>™</sup>  
REAL INFORMATION FROM REAL PATIENTS  
Only from Sandostatin® LAR Depot  
(octreotide acetate) for injectable suspension  
10mg • 20mg • 30mg

For more recipes, tips, and helpful resources,  
like us on [Facebook.com/SandostatinLARDepot](https://www.facebook.com/SandostatinLARDepot)

*This recipe was created by our RealSupport™ nutritionist especially for our members. Remember, please talk to your health care provider to determine if this recipe or any other food is appropriate for you.*

# RICOTTA SCRAMBLED EGG AND ASPARAGUS TOAST

 Prep time: 5 minutes

 Cook time: 10 minutes

 Ready in: 15 minutes

## Ingredients

- 2 tablespoons olive oil
- ½ teaspoon kosher salt, divided
- ¼ teaspoon black pepper, plus more for serving
- 4 large eggs
- 1 teaspoon chives, finely chopped
- 1 tablespoon unsalted butter
- ½ cup fresh low-fat ricotta
- ½ pound asparagus, ends trimmed
- 2 large (or 4 small) thick slices rustic white bread, lightly toasted

## Steps

1. Heat oil in large nonstick skillet over medium-high heat. Add asparagus, ¼ teaspoon kosher salt, and black pepper, stirring to coat. Cook for 10 minutes, stirring frequently to ensure even cooking.
2. While asparagus cooks, prepare eggs. Whisk eggs, chives, and remaining salt together until well-beaten. Melt unsalted butter in a medium nonstick skillet over medium heat.
3. Add eggs to melted butter and slowly stir with silicone spatula, scraping from the edges toward the center. Continue stirring until eggs are mostly cooked but still soft, about 2 minutes. Remove from heat and stir in low-fat ricotta.
4. Spoon eggs over toast and top with cooked asparagus spears. Garnish with black pepper.

\*Includes Sandostatin® (octreotide acetate) Immediate-Release Injection and Sandostatin® LAR Depot (octreotide acetate) for injectable suspension for all approved indications.

**Please note that Novartis Pharmaceuticals Corporation does not endorse the use of any foods highlighted in this article; all information is provided for educational purposes only. Use of any food in this recipe is not intended to treat or cure carcinoid syndrome. You are advised to speak to your health care provider before introducing new foods into your diet.**

**Please see Indications and Important Safety Information on page 2.**

Please [click here](#) for full Prescribing Information.

Brought to you by RealSupport™, a patient information and support program for those with the severe diarrhea and flushing of carcinoid syndrome. RealSupport™ is provided by Novartis Pharmaceuticals Corporation, the maker of Sandostatin® LAR Depot (octreotide acetate) for injectable suspension. For more information about RealSupport™ program offerings, please follow us on [www.facebook.com /SandostatinLARDepot](https://www.facebook.com/SandostatinLARDepot).

## INDICATIONS AND USAGE

Sandostatin® LAR Depot (octreotide acetate) for injectable suspension is indicated for patients in whom initial treatment with immediate-release Sandostatin® (octreotide acetate) Injection has been shown to be effective and tolerated for

- Long-term treatment of the severe diarrhea and flushing episodes associated with metastatic carcinoid tumors
- Long-term treatment of the profuse watery diarrhea associated with VIP-secreting tumors

In patients with carcinoid syndrome and VIPomas, the effect of Sandostatin Injection and Sandostatin LAR Depot on tumor size, rate of growth, and development of metastases has not been determined.

## IMPORTANT SAFETY INFORMATION

**Warnings and Precautions:** Treatment with Sandostatin LAR Depot may affect gallbladder function, with postmarketing reports of gallstones resulting in complications (inflammation of the gallbladder, bile duct, and pancreas, and requiring surgical removal of the gallbladder); sugar metabolism; thyroid and heart function; and nutritional absorption, which may require monitoring by your doctor. Call your doctor if you experience signs or symptoms of gallstones or any of their complications.

**Before Taking Sandostatin LAR Depot:** Tell your doctor if you have a history of heart disease or are taking other medications, including cyclosporine, insulin, oral hypoglycemic agents, beta-blockers, and bromocriptine.

**Common Side Effects:** Most patients experience side effects at some time. Some common side effects you may experience include back pain, fatigue, headache, abdominal pain, nausea, and dizziness.

**Other Information:** Patients with carcinoid tumors and VIPomas should adhere closely to their scheduled return visits for reinjection in order to minimize exacerbation of symptoms.

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.**

Please [click here](#) for full Prescribing Information.

To learn more, call 1-888-NOW-NOVA (1-888-669-6682) or visit [us.sandostatin.com](http://us.sandostatin.com)

